




## Restriction Enzyme Eco47 I



Cat.#	Size	Conc.
FG-Eco47I	800 units	10 units/μl

Store at -20°C

**Supplied with:** 10X FastGene® Buffer III (FG-REB3)  
10X FastGene® FastCut Buffer (FG-REBHF)  
6X DNA Loading Buffer  
Sterile water

### Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

**Source:** *E.coli* RFL47

### Reaction conditions

1X FastGene® Buffer III 37°C  
1X FastGene® FastCut Buffer, 37°C

### FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

### 1X FastGene® Buffer III

50 mM Tris-HCl (pH 7.9 at 25°C)  
100 mM NaCl  
10 mM MgCl<sub>2</sub>  
100 μg/ml BSA

### Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ at 37°C for 1 hr in 50 μl reaction mixtures.

### Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

### Dilution buffer:

FastGene® Diluent A

### Heat Inactivation

Eco47 I can be inactivated at 65°C for 20 min.

### Methylation sensitivity

*dam* methylation: Not sensitive  
*dcm* methylation: sensitive  
CpG methylation: sensitive

### Relative activity in FastGene® Buffers

FastGene® Buffer I: 100%  
FastGene® Buffer II: 100%  
FastGene® Buffer III: 100%  
FastGene® Buffer IV: 100%  
FastGene® FastCut Buffer: 100%

### Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer III	1 X	5 μl
Eco47 I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® FastCut Buffer	1 X	5 μl
Eco47 I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.



## Restriction Enzyme Eco47 I



Cat.#	Size	Conc.
FG-Eco47I	800 units	10 units/μl

Store at -20°C

**Supplied with:** 10X FastGene® Buffer III (FG-REB3)  
10X FastGene® FastCut Buffer (FG-REBHF)  
6X DNA Loading Buffer  
Sterile water

### Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

**Source:** *E.coli* RFL47

### Reaction conditions

1X FastGene® Buffer III 37°C  
1X FastGene® FastCut Buffer, 37°C

### FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

### 1X FastGene® Buffer III

50 mM Tris-HCl (pH 7.9 at 25°C)  
100 mM NaCl  
10 mM MgCl<sub>2</sub>  
100 μg/ml BSA

### Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ at 37°C for 1 hr in 50 μl reaction mixtures.

### Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

### Dilution buffer:

FastGene® Diluent A

### Heat Inactivation

Eco47 I can be inactivated at 65°C for 20 min.

### Methylation sensitivity

*dam* methylation: Not sensitive  
*dcm* methylation: sensitive  
CpG methylation: sensitive

### Relative activity in FastGene® Buffers

FastGene® Buffer I: 100%  
FastGene® Buffer II: 100%  
FastGene® Buffer III: 100%  
FastGene® Buffer IV: 100%  
FastGene® FastCut Buffer: 100%

### Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer III	1 X	5 μl
Eco47 I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® FastCut Buffer	1 X	5 μl
Eco47 I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.